普诺赛[®] Procell system

VEGF-A/VEGF121, Human, Recombinant

Cat. No.: GPCK013

产品信息

物种 Human

表达宿主 E.coli

序列信息 Ala27-Arg147 检索号 P15692-9 分子量 14.2 kDa 有效期 12 months

生物活性 Immobilized Human VEGF 121 at 2 µg/mL (100 µL/well) can bind Human VEGFR2-Fc. 隊® | Procell system

cell system

The ED50 of Human VEGFR2-Fc is 13.04 ng/mL.

产品特性

内毒素 (EU/μg) < 0.1

保存 Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt.

Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at -5~-20°C for 3 months.

运输 Ambient temperature or ice pack.

制剂 Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

复溶 Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not

> recommended to reconstitute to a concentration less than 100 µg/mL. Dissolve the lyophilized protein in sterile water. Please aliquot the reconstituted solution to minimize

freeze-thaw cycles.

背景介绍

Human VEGF121, also known as Vascular Endothelial Growth Factor A, VEGFA, Vascular Permeability Factor, VPF and VEGF, is a homodimeric, heparin-binding glyco Protein which belongs to the platelet-Derived Growth Factor (PDGF)/vascular Endothelial Growth Factor (VEGF) family. VEGF-A is a glycosylated mitogen that specifically acts on Endothelial cells and has various effects, including mediating increased vascular Permeability, inducing angiogenesis, vasculogenesis, permeabilization of blood vessels and Endothelial cell Growth, increasing microvascular Permeability, promoting cell migration and inhibiting apoptosis. Alternatively spliced transcript variants of VEGF-A encod either secreted or cell-associated isoforms. The lymphangiogenesis may be promoted by upregulation of VEGF121, which may in turn act in part via induction of VEGF-C. It binds to the FLT1/VEGFR1 and KDR/VEGFR2 Receptors, heparan sulfate and heparin. NRP1/Neuropilin-1 binds isoforms VEGF-165 and VEGF-145. Isoform VEGF165B binds to KDR but does not activate downstream signaling pathways, does not activate angiogenesis and inhibits tumor Growth.

网站: www.procell.com.cn 电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋



