

IL-13/NC30, Human, Recombinant**Cat. No. : GPCK362****产品信息**

物种	Human
表达宿主	E.coli
序列信息	MGPVPPSTALRELIEELVNITQNQKAPLCNGSMVWSINLTAGMYCAALESLINVSGCSAIEKTQ RMLSGFCPHKVSAGQFSSLHVRDTKIEVAQFVKDLLHLKKLRFREGRFN with polyhistidine tag at the C-terminus
检索号	P35225.2
标签	His-tag at the C-terminus
分子量	13.28 kDa
有效期	12 months
生物活性	Measure by its ability to induce TF-1 cells proliferation. The ED50 for this effect is < 0.8 ng/mL. The specific activity of recombinant human IL-13 is approximately > 1 × 10 ⁶ IU/ mg.

产品特性

内毒素 (EU/μg)	< 0.1
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 μm filtered solution containing 1 × PBS, pH 8.0.
复溶	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 μg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

背景介绍

Interleukin 13 (IL-13) is a protein that in humans is encoded by the IL13 gene. IL-13 was first cloned in 1993 and is located on chromosome 5q31 with a length of 1.4kb. It has a mass of 13 kDa and folds into 4 alpha helical bundles. The secondary structural features of IL-13 are similar to that of Interleukin 4 (IL-4); however it only has 25% sequence homology to IL-4 and is capable of IL-4 independent signaling. IL-13 is a cytokine secreted by T helper type 2 (Th2) cells, CD4 cells, Natural killer T cell, Mast cell, Basophil cells, Eosinophil cells and Nuocyte cells. Interleukin-13 is a central regulator in IgE synthesis, goblet cell hyperplasia, mucus hypersecretion, airway hyperresponsiveness, fibrosis and chitinase up-regulation. It is a mediator of allergic inflammation and different diseases including asthma.

网站: www.procell.com.cn

电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋



普诺赛® | Procell system

普诺赛® | Procell system

普诺赛® | Procell system

普诺赛® | Procell system

