

PE Anti-Human CD160 Antibody[BY55]

Catalog Number: AN01145D

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity	Human
Host	Mouse
Isotype	Mouse IgM, κ
Clone No.	BY55
Isotype Control	PE Mouse IgM, κ Isotype Control[MM-30] [Product E-AB-F09782D]
Conjugation	PE
Conjugation Information	PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green (561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.


Applications

Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of normal human peripheral blood cells with APC Anti-Human CD19 Antibody and PE Anti-Human CD160 Antibody[BY55] (left) or PE Mouse IgM, κ Isotype Control (right). Cells in the lymphocytes gate were used for analysis.

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	BY55CNK1;NK28
Uniprot ID	O95971
Gene ID	11126

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.3

Background

CD160 is a 27 kD GPI-anchored glycoprotein also known as BY55, NK1, and NK28. A member the Ig superfamily, CD160 exists as a disulfide-bond multimer, expressed on the surface of a subpopulation of NK cells, $\gamma\delta$ T cells, subset of CD8+ T cells, and intestinal intraepithelial lymphocytes (IEL). CD160 plays costimulatory roles through binding to classical and nonclassical MHC-I molecules.