Recombinant U2AF2 Monoclonal Antibody

catalog number: **AN302058L**

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description

Reactivity	Human;Mouse;Rat
Immunogen	Peptide. This information is proprietary to PTMab.
Host	Rabbit
Isotype	Rabbit IgG
Clone	C121679
Purification	Protein A purified
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications	Recommended Dilution
WB	1:5000-1:10000
IHC	1:200
ChIP	6 μg/ 2×10^6 cells

Data



Western Blot with U2AF2 Monoclonal Antibody at dilution of 1:10000. Lane 1: HEK-293, Lane 2: HeLa, Lane 3: Jurkat, Lane 4: NIH-3T3

Observed-MW:65 kDa

Calculated-MW:54 kDa



Immunohistochemistry of paraffin-embedded Human colon cancer using U2AF2 Monoclonal Antibody at dilution of 1:200.



Immunohistochemistry of paraffin-embedded Mouse liver using U2AF2 Monoclonal Antibody at dilution of 1:200.



Immunohistochemistry of paraffin-embedded Rat cerebrum using U2AF2 Monoclonal Antibody at dilution of 1:200.



Chromatin immunoprecipitation analysis of HepG2 immunoprecipitated DNA by real-time PCR using primers specific for the human ZNF446-E and PPCS-I. The data are presented as enrichment of each sample relative to the total amount of input chromatin at each amplicon.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

TFIIF is a general transcription initiation factor that binds to RNA polymerase II and helps to recruit it to the initiation complex in collaborationwith TFIIB.

Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web: www.elabscience.com
Email: techsupport@elabscience.com