


Recombinant U2AF2 Monoclonal Antibody

catalog number: **AN302058L**


Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description	
Reactivity	Human;Rat;Mouse
Immunogen	Peptide. This information is proprietary to PTMap.
Host	Rabbit
Isotype	IgG, κ
Clone	C121679
Purification	Protein A purified
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.
Applications	
WB	1:5000-1:10000
IHC	1:200
ChIP	6 μg/ 2×10 ⁶ cells
Data	




Western Blot with U2AF2 Monoclonal Antibody at dilution of 1:10000. Lane 1: HEK-293, Lane 2: HeLa, Lane 3: Jurkat, Lane 4: NIH-3T3


Observed-MW:65 kDa
Calculated-MW:54 kDa




Immunohistochemistry of paraffin-embedded Human colon cancer using U2AF2 Monoclonal Antibody at dilution of 1:200.



Immunohistochemistry of paraffin-embedded Mouse liver using U2AF2 Monoclonal Antibody at dilution of 1:200.



Immunohistochemistry of paraffin-embedded Rat cerebrum using U2AF2 Monoclonal Antibody at dilution of 1:200.



For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.2

Chromatin immunoprecipitation analysis of HepG2 immunoprecipitated DNA by real-time PCR using primers specific for the human ZNF446-E and PPCS-I. The data are presented as enrichment of each sample relative to the total amount of input chromatin at each amplicon.

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping Ice bag

Background

TFIIF is a general transcription initiation factor that binds to RNA polymerase II and helps to recruit it to the initiation complex in collaboration with TFIIB.