## Elabscience Biotechnology Co., Ltd.



A Reliable Research Partner in Life Science and Medicine

# Recombinant Cynomolgus CD40/TNFRSF5/CD40L receptor Protein (His Tag)

Catalog Number: PKSQ050045

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

**Species** Cynomolgus macaques

Source HEK293 Cells-derived Cynomolgus macaques CD40/TNFRSF5 protein Glu21-Arg 193,

with an C-terminal His

Calculated MW20.2 kDaObserved MW28-30 kDaAccessionG7PG38

**Bio-activity** Not validated for activity

## **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin**  $< 1.0 \text{ EU per } \mu\text{g of the protein as determined by the LAL method.}$ 

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

#### Background

CD40 is a Type I Transmembrane Glycoprotein that belongs to the TNF Receptor Superfamily. CD40 is expressed in B cells, follicular dendritic cells, dendritic cells, activated monocytes, macrophages, endothelial cells, vascular smooth muscle cells, and several tumor cell lines. The extracellular domain of CD40 is characterized by Cysteine rich repeat regions. Interaction of CD40 with its ligand (CD40L) leads to aggregation of CD40 molecules, which in turn interact with cytoplasmic components to initiate signaling pathways. Several different TRAF proteins (adaptor proteins) have been identified to serves as mediators of the signal transduction.

#### For Research Use Only