

TAF5 Polyclonal Antibody

catalog number: E-AB-61593

Note: Centrifuge before opening to ensure complete recovery of vial contents.



Description

Reactivity	Human;Mouse;Rat
Immunogen	Recombinant fusion protein of human TAF5 (NP_008882.2).
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications

Applications	Recommended Dilution
WB	1:500-1:2000
IF	1:50-1:100

Data



Western blot analysis of extracts of Mouse brain using TAF5 Polyclonal Antibody at dilution of 1:1000.

Immunofluorescence analysis of A-549 cells using TAF5 Polyclonal Antibody

Observed-MW: 100 kDa

Calculated-MW: 80 kDa/86 kDa

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes an integral subunit of TFIID associated with all transcriptionally competent forms of that complex. This subunit interacts strongly with two TFIID subunits that show similarity to histones H3 and H4, and it may participate in forming a nucleosome-like core in the TFIID complex. Alternative splicing results in multiple transcript variants.

For Research Use Only