A Reliable Research Partner in Life Science and Medicine

Recombinant Mouse GFRA1/GDNFRA Protein (His Tag)

Catalog Number: PKSM040827

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species Mouse

Source HEK293 Cells-derived Mouse GFRA 1/GDNFRA protein Met 1-Ser 425, with an C-

terminal His

Calculated MW 46.3 kDa Observed MW 55-60 kDa Accession AAB86600.1

Bio-activity 1. Measured in a cell proliferation assay using SH- SY5Y human neuroblastoma cells.

> The ED $_{50}$ for this effect is typically 0.3-1. 5 $\mu g/mL$ in the presence of 40 ng/mLRecombinant Human GDNF. 2. Measured by its ability to bind with human GDNF in a

functional ELISA.

Properties

> 98 % as determined by reducing SDS-PAGE. **Purity**

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 Storage

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

This product is provided as lyophilized powder which is shipped with ice packs. Shipping

Lyophilized from sterile PBS, pH 7.4 **Formulation**


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Elabscience®

Glial cell line derived neurotrophic factor (GDNF) Family Receptor Alpha 1 (GFRA1) is a member of the GDNF receptor family. It is a glycosylphosphatidylinositol (GPI)-linked cell surface receptor for both GDNF and NTN, and mediates activation of the RET tyrosine kinase receptor. GFRA1 is a potent survival factor for central and peripheral neurons, and is essential for the development of kidneys and the enteric nervous system. Glial cell line-derived neurotrophic factor (GDNF) and neurturin (NTN) are its binding ligand which are two structurally related, potent neurotrophic factors that play key roles in the control of neuron survival and differentiation. GDNF promotes the formation of a physical complex between GFRA/GDNFRa and the orphan tyrosin kinase receptor Ret, thereby inducing its tyrosine phosphorylation. The RET is a receptor tyrosine kinase representing the signal-transducing molecule of a multisubunit surface receptor complex for the GDNF, in which GFRA / GDNFRa acts as the ligand-binding component. GDNF, a distantly related member of the transforming growth factor-β (TGF-â) superfamily, and its receptor components: GFRA1, Ret and neural cell adhesion molecule (NCAM) have been recently reported to be expressed in the testis and to be involved in the proliferation regulation of immature Sertoli cells.

Toll-free: 1-888-852-8623 Web:www.elabscience.com Fax: 1-832-243-6017