Recombinant Human PLAUR Protein(His Tag)

Catalog Number: PDMH100351

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Human
Source	Mammalian-derived Human PLAUR proteins Met1-Arg303, with an C-terminal His
Calculated MW	33.2 kDa
Observed MW	45 kDa
Accession	Q03405
Bio-activity	Not validated for activity
Properties	
Purity	>90% as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS with 5% Trehalose and 5%
	Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of
	0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Human PLAUR proteins, 2 µg/lane of Recombinant Human PLAUR proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 45

KD

Background

Elabscience®

Urokinase plasminogen activator (uPA) and/or its receptor (uPAR) are essential for metastasis, and overexpression of these molecules is strongly correlated with poor prognosis in a variety of malignant tumours. uPAR and uPA levels in both resected tumor tissue and plasma are of independent prognostic significance for patient survival in several types of human cancer. This system has classically been thought to drive tumor progression by mediating directed extracellular proteolysis on the surface of migrating or invading cells, and intervening with this proteolysis by targeting uPAR has been proposed to represent a novel approach for inhibiting tumor progression. uPAR, also known as PLAUR or CD87, has been implicated in the growth, metastasis, and angiogenesis of several solid and hemotologic malignancies. uPAR is a highly glycosylated, 55-60 KD integral membrane protein linked to the plasma membrane by a glycosylphosphatidylinositol (GPI) anchor. It is part of a cell surface system that also consists of the serine protease uPA and several specific inhibitors (plasminogen activator inhibitors 1 and 2). Additionally, the analysis of CD87 (urokinase-type plasminogen activator receptor-uPAR) expression has a potential role in the diagnostic or prognostic work-up of several hematological malignancies, particularly acute leukemia and multiple myeloma.