

SNRPA Polyclonal Antibody

catalog number: E-AB-61942

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse; Rat

Immunogen Recombinant fusion protein of human SNRPA (NP 004587.1).


Host Rabbit Isotype IgG


Purification Affinity purification

Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications	Recommended Dilution	
IHC	1:50-1:200	
IF	1:50-1:200	


Data







Immunohistochemistry of paraffin-embedded Rat liver using Immunohistochemistry of paraffin-embedded Rat brain using SNRPA Polyclonal Antibody at dilution of 1:100 (40x lens). SNRPA Polyclonal Antibody at dilution of 1:100 (40x lens).





Immunohistochemistry of paraffin-embedded Rat heart using Immunohistochemistry of paraffin-embedded Human breast SNRPA Polyclonal Antibody at dilution of 1:100 (40x lens). using SNRPA Polyclonal Antibody at dilution of 1:100 (40x lens).





Immunohistochemistry of paraffin-embedded Human stomach using SNRPA Polyclonal Antibody at dilution of 1:100 (40x lens).


Immunohistochemistry of paraffin-embedded Human gastric cancer using SNRPA Polyclonal Antibody at dilution of 1:100 (40x lens).

For Research Use Only

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine



Immunofluorescence analysis of HeLa cells using SNRPA
Polyclonal Antibody

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

The protein encoded by this gene associates with stem loop II of the U1 small nuclear ribonucleoprotein, which binds the 5' splice site of precursor mRNAs and is required for splicing. The encoded protein autoregulates itself by polyadenylation inhibition of its own pre-mRNA via dimerization and has been implicated in the coupling of splicing and polyadenylation.

Fax: 1-832-243-6017