Recombinant Human ANPEP Protein(His Tag)


Note: Centrifuge before opening to ensure complete recovery of vial contents.

Catalog Number: PDMH100269



Description Species Human Source Mammalian-derived Human ANPEP protein Lys69-Lys967, with an C-terminal His Mol Mass 98.8 kDa P15144 Accession **Bio-activity** Not validated for activity **Properties** >90% as determined by reducing SDS-PAGE. Purity Endotoxin < 1.0 EU/mg of the protein as determined by the LAL method Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 Storage °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at $< -20^{\circ}$ C for 3 months. Shipping This product is provided as lyophilized powder which is shipped with ice packs. Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Formulation Mannitol. Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Human ANPEP proteins, 2 µg/lane of Recombinant Human ANPEP proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 98.8 KD

Background

Aminopeptidase N (ANPEP or APN), also known as CD13, is a cell-surface metalloprotease located in the small-intestinal and renal microvillar membrane, as well as other plasma membranes. It belongs to the peptidase M1 family. CD13 plays a role in the final digestion of peptides generated from hydrolysis of proteins by gastric and pancreatic proteases and is involved in the metabolism of regulatory peptides by diverse cell types. CD13/APN is a potent regulator of angiogenesis which is essential for tumor invasion and metastasis, and its transcription in activated endothelial cells is induced by angiogenic growth factors via the RAS/MAPK pathway. In addition, this enzyme has been shown to participate in antigen processing and presentation, and accordingly, defects in this gene appear to be a cause of various types of leukemia or lymphoma and carcinomas.

For Research Use Only