

PE/Cyanine5 Anti-Human TCR Va24-Ja18 Antibody[6B11]

Catalog Number: AN00568G

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity	Human
Host	Mouse
Isotype	Mouse IgG1
Clone No.	6B11
Isotype Control	PE/Cyanine5 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792G]
Conjugation	PE/Cyanine 5
Conjugation Information	PE/Cyanine5 is designed to be excited by the Blue (488 nm), Green (532 nm) and yellow-green (561 nm) lasers and detected using an optical filter centered near 670 nm (e.g., a 690/50 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.


Applications

Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of normal human peripheral blood cells with FITC Anti-Human CD3 Antibody and PE/Cyanine5 Anti-Human TCR Va24-Ja18 Antibody[6B11] (left) or PE/Cyanine5 Mouse IgG1, κ Isotype Control (right). Cells in the lymphocytes gate were used for analysis.

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	TCR Va24-Ja18;TCR Va24-JaQ;invariant NKT cell;iNKT
Gene ID	28659

For Research Use Only

Background

Encoded by the TCR Va24-Ja18 germline configuration, Va24-JaQ is expressed on a subset of NKT cells, namely invariant NKT (iNKT). Va24-JaQ TCR interacts with the glycolipid loaded MHC class 1b molecule CD1d, inducing activation and subsequent cytokine production. iNKT cells have been implicated in immune regulation, tumor surveillance, and host response to pathogens. While iNKT cells occur at low frequency in the blood, assorted chemokines contribute to their tissue homing potential.