

## PE Anti-Mouse NKG2A/C/E Antibody[20d5]

Catalog Number: AN00409D

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


<b>Reactivity</b>	Mouse
<b>Host</b>	Rat
<b>Isotype</b>	Rat IgG2a, κ
<b>Clone No.</b>	20d5
<b>Isotype Control</b>	PE Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832D]
<b>Conjugation</b>	PE
<b>Conjugation Information</b>	PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green (561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42 nm bandpass filter).
<b>Storage Buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

### Applications

### Recommended usage

<b>FCM</b>	Each lot of this antibody is quality control tested by flow cytometric analysis. <b>The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood).</b> Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
------------	---

### Data



Staining of C57BL/6 murine splenocytes cells with APC Anti-Mouse NK1.1 Antibody and FITC Anti-Mouse NKG2A/C/E Antibody[20d5] (left) or FITC Rat IgG2a,κ Isotype Control (right). Total viable cells were used for analysis.

### Preparation & Storage

<b>Storage</b>	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze.
<b>Shipping</b>	Ice bag

### Antigen Information

<b>Alternate Names</b>	KLRC1;Killer Cell Lectin Like Receptor C1;KLRC2;Killer Cell Lectin Like Receptor C2; KLRC3;Killer Cell Lectin Like Receptor C3
<b>Uniprot ID</b>	P26715

### For Research Use Only

**Gene ID**

500338

**Background**

The NKG2 molecules are a family of lectin-like receptors that form heterodimers with CD94. NKG2/CD94 heterodimer are primarily expressed on NK cells, and a subset of CD8+ T cells. Studies of CD94/NKG2 heterodimers on NK cells have demonstrated that the NKG2 components transduce signals after ligand binding. NKG2A transduces inhibitory signals, while NKG2C and NKG2E transduce activating signals.