

PE Anti-Mouse VISTA Antibody[MIH63]

Catalog Number: AN00875D

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Mouse
Host	Rat
Isotype	Rat IgG2a
Clone No.	MIH63
Isotype Control	PE Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832D]
Conjugation	PE
Conjugation Information	PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green (561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications

Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of C57BL/6 murine splenocytes with Elab Fluor® Violet 450 Anti-Mouse CD8a Antibody[53-6.7] and PE Anti-Mouse VISTA Antibody[MIH63](left) or PE Rat IgG2a, κ Isotype Control (right). Total viable cells were used for analysis.

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protect from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	PD-1 homolog; Dies1; B7-H5
Uniprot ID	Q9D659

For Research Use Only

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.3

Gene ID	74048
Background	PD-1H, also known as VISTA, is a 309 aa type I transmembrane protein that is composed of seven exons. PD-1H has one Ig-V like domain, and its sequence is similar to the Ig-V domains of the members of CD28 and B7 families. PD-1H is expressed by a subset of T cells, macrophages, dendritic cells, neutrophils, and natural killer cells (NK). It has been proposed that PD-1H can be useful to modulate the host immune response to allogeneic transplants due to its ability to preferentially suppress CD4+ T cell-mediated immunity.

For Research Use Only

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.3