







MTHFD2 Polyclonal Antibody

catalog number: E-AB-65766

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description	
Reactivity	Human;Mouse;Rat
Immunogen	Recombinant fusion protein of human MTHFD2 (NP_006627.2).
Host	Rabbit
Isotype	Rabbit IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
IHC	1:50-1:200
IF	1:50-1:200

Data	
 Immunohistochemistry of paraffin-embedded Human mammary cancer using MTHFD2 Polyclonal Antibody at dilution of 1:100 (40x lens).	 Immunohistochemistry of paraffin-embedded Human liver cancer using MTHFD2 Polyclonal Antibody at dilution of 1:100 (40x lens).
 Immunohistochemistry of paraffin-embedded Rat kidney using MTHFD2 Polyclonal Antibody at dilution of 1:100 (40x lens).	 Immunohistochemistry of paraffin-embedded Mouse kidney using MTHFD2 Polyclonal Antibody at dilution of 1:100 (40x lens).
 Immunofluorescence analysis of C6 cells using MTHFD2 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.	 Immunofluorescence analysis of NIH-3T3 cells using MTHFD2 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.


For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.7



Immunofluorescence analysis of U-2 OS cells using MTHFD2 Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

This gene encodes a nuclear-encoded mitochondrial bifunctional enzyme with methylenetetrahydrofolate dehydrogenase and methenyltetrahydrofolate cyclohydrolase activities. The enzyme functions as a homodimer and is unique in its absolute requirement for magnesium and inorganic phosphate. Formation of the enzyme-magnesium complex allows binding of NAD. Alternative splicing results in two different transcripts, one protein-coding and the other not protein-coding. This gene has a pseudogene on chromosome 7.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.7