A Reliable Research Partner in Life Science and Medicine

AF/LE Purified Anti-Human CD86 Antibody[IT2.2]

catalog number: E-AB-F12690

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Recombinant Human CD86 protein

Host Mouse

Isotype Mouse IgG2b, κ

Clone IT2.2


Purification >98%, Protein A/G purified

Buffer Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method

Applications Recommended Dilution

FCM $2 \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$

Data



Human peripheral blood monocytes cell were stained with 0.2 μ g AF/LE Purified Anti-Human CD86 Antibody[IT2.2] (Right) and 0.2 μ g Mouse IgG2b, κ Isotype Control (Left), followed by Alexa Fluor® 647-conjugated Goat Anti-Mouse

IgG Secondary Antibody.

Preparation & Storage

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles. This preparation contains no preservatives, thus it should be handled

under aseptic conditions.

Shipping Ice bag

Background

This gene encodes a type I membrane protein that is a member of the immunoglobulin superfamily. This protein is expressed by antigen-presenting cells, and it is the ligand for two proteins at the cell surface of T cells, CD28 antigen and cytotoxic T-lymphocyte-associated protein 4. Binding of this protein with CD28 antigen is a costimulatory signal for activation of the T-cell. Binding of this protein with cytotoxic T-lymphocyte-associated protein 4 negatively regulates T-cell activation and diminishes the immune response. Alternative splicing results in several transcript variants encoding different isoforms.

For Research Use Only

Fax: 1-832-243-6017