A Reliable Research Partner in Life Science and Medicine

Recombinant Human Semaphorin-4D/CD100 protein (His Tag)

Catalog Number: PDMH100385

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species Human

Source HEK293 Cells-derived Human SEMA4D protein Met1-Arg734, with an C-terminal His

 Calculated MW
 80.6 kDa

 Observed MW
 90-110 kDa

 Accession
 Q92854

Bio-activity Not validated for activity

Properties

Purity > 95% as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU/mg of the protein as determined by the LAL method

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.


ShippingThis product is provided as lyophilized powder which is shipped with ice packs. **Formulation**Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5%

Mannitol.

Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution of

0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Human Semaphorin-4D/CD100 proteins, 2µg/lane of Recombinant Human Semaphorin-4D/CD100 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 90-110

KD.

Background

Elabscience®

Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

Semaphorin-4D is also known as A8,BB18, GR3, CD100. Semaphorin-4D belongs to the semaphorin family containing 1 Ig-like C2-type domain, 1 PSI domain and 1 Sema domain. It is the cell surface receptor for PLXN1B and PLXNB2 that plays an important role in cell-cell signaling. It promotes the migration of cerebellar granule cells and of endothelial cells, regulates dendrite and axon branching and morphogenesis. Semaphorin-4D Plays a role in the immune system, Promotes signaling via SRC and PTK2B/PYK2, which then mediates activation of phosphatidylinositol 3-kinase and of the AKT1 signaling cascade.