Elabscience Biotechnology Co., Ltd.



A Reliable Research Partner in Life Science and Medicine

RAF1 Polyclonal Antibody

catalog number: E-AB-60020

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse; Rat

Immunogen Recombinant fusion protein of human RAF1 (NP 002871.1).

Host Rabbit Isotype IgG


Purification Affinity purification
Conjugation Unconjugated

Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.


Applications	Recommended Dilution
WR	1:500-1:2000

HC 1:50-1:200 IF 1:50-1:200

Data




Western blot analysis of extracts of various cell lines using RAF1 Polyclonal Antibody at dilution of 1:1000.




Immunohistochemistry of paraffin-embedded Rat liver using RAF1 Polyclonal Antibody at dilution of 1:100 (40x lens).

Observed-MW:73 kDa Calculated-MW:73 kDa/75 kDa



Immunohistochemistry of paraffin-embedded Rat heart using RAF1 Polyclonal Antibody at dilution of 1:100 (40x lens).



Immunofluorescence analysis of U2OS cells using RAF1
Polyclonal Antibody

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

Web: www.elabscience.cn

temperature recommended.

Background

For Research Use Only



Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

This gene is the cellular homolog of viral raf gene (v-raf). The encoded protein is a MAP kinase kinase kinase (MAP3K), which functions downstream of the Ras family of membrane associated GTPases to which it binds directly. Once activated, the cellular RAF1 protein can phosphorylate to activate the dual specificity protein kinases MEK1 and MEK2, which in turn phosphorylate to activate the serine/threonine specific protein kinases, ERK1 and ERK2. Activated ERKs are pleiotropic effectors of cell physiology and play an important role in the control of gene expression involved in the cell division cycle, apoptosis, cell differentiation and cell migration. Mutations in this gene are associated with Noonan syndrome 5 and LEOPARD syndrome 2.

Web: www.elabscience.cn