## **Elabscience**®

## **ERK 1 Monoclonal Antibody**

## catalog number: E-AB-22246

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description                  |  |   |  |
|------------------------------|--|---|--|
| Reactivity                   | Human;Mouse;Rat  |   |  |
| •                            |  | Recombinant Protein of ERK1 of MAPK3  |  |
| Immunogen                    |  |   |  |
| Host                         | Mouse  |   |  |
| Isotype                      | IgG  | -   |  |
| Clone                        | 1C11   |   |  |
| Purification                 | Protein A purification   | •   |  |
| Conjugation                  | Unconjugated   |   |  |
| Buffer                       | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 0.5% protein |   |  |
| protectant and 50% glycerol. |  | erol.   |  |
| Applications                 | <b>Recommended Dilution</b>  | Recommended Dilution  |  |
| IHC                          | 1:100-200  |   |  |
| Data                         |  |   |  |
|                              |  |   |  |
| Immunohistochemistry of p    | paraffin-embedded Human tonsil   | Immunohistochemistry of paraffin-embedded Human breast                          |  |
| tissue using ERK 1 Mono      | oclonal Antibody at dilution of  | carcinoma tissue using ERK 1 Monoclonal Antibody at                             |  |
|                              | 1:200.   | dilution of 1:200.  |  |
| Preparation & Storage        |  |   |  |
| Storage                      | Store at -20°C Valid for 12  | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.                 |  |
| Shipping                     | The product is shipped w   | The product is shipped with ice pack, upon receipt, store it immediately at the |  |
|                              | temperature recommended.   |   |  |
| Background                   |  |   |  |

Involved in both the initiation and regulation of meiosis, mitosis, and postmitotic functions in differentiated cells by phosphorylating a number of transcription factors such as ELK-1. Phosphorylates EIF4EBP1; required for initiation of translation. Phosphorylates microtubule-associated protein 2 (MAP2). Phosphorylates SPZ1 (By similarity). Phosphorylates heat shock factor protein 4 (HSF4).