



A Reliable Research Partner in Life Science and Medicine

Recombinant Human IL-30 protein(N-His)

Catalog Number: PKSH034115

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| - | | | | | 70 | . • | | |
|-----|----|---|----|---|----|-----|---|--|
| H)) | es | C | r٦ | n | tı | n | m | |

Species Human

Source E.coli-derived Human IL-30 protein Phe 29-Pro 243, with an C-terminal His

 Mol_Mass
 25.5 kDa

 Accession
 Q8NEV9

Bio-activity Not validated for activity

Properties

Purity > 98 % as determined by reducing SDS-PAGE.

Endotoxin < 0.1 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS,pH 8.0.

Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Background

IL-30, also known as IL-27A,p28. IL-27 protein is a member of the IL-6 superfamily, which is expressed on monocytes, endothelial cells, and dendritic cells. IL-27 protein is also referred to as the IL-12 p35-related protein, p28, is one subunit of a heterodimeric cytokine complex, and associates with another subunit EBI3 (EBV-induced gene 3), and IL-12 p40-related protein (IL-27B). IL-27 protein is an early product of activated antigen-presenting cells and drives the rapid clonal expansion of naive CD4(+) T cells and plays a role in the early regulation of Th1 cells initiation which drives efficient adaptive immune response. IL-27 protein has an antiproliferative activity on melanomas through WSX-1/STAT1 signaling. Thus, IL-27 protein may be an attractive candidate as an antitumor agent applicable to cancer immunotherapy.