

Recombinant Human Receptor Tyrosine-Protein Kinase ErbB-2/HER2 (C-Fc-Avi) Biotinylated

Catalog Number: PKS033959

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human HER2 protein Thr23-Thr652, with an C-terminal Fc & Avi
Calculated MW	98.3 kDa
Observed MW	120-140 kDa
Accession	P04626
Bio-activity	Not validated for activity

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Human epidermal growth factor receptor 2 (HER2) is a type of membrane glycoprotein, and belongs to the epidermal growth factor (EGF) receptor family. HER2 plays a key role in development, cell proliferation and differentiation. HER2 has been reported to associate with malignancy and a poor prognosis in numerous carcinomas, including breast, prostate, ovarian, lung cancers and so on. HER2 is activated by dimerization and not activated by EGF, TGF- α and amphiregulin. Interaction with PTK6 increases its intrinsic kinase activity. It is heterodimer with EGFR, ERBB3 and ERBB4. HER2 associates with the 5'-TCAAATTC-3' sequence in the PTGS2/COX-2 promoter and activates its transcription. It is implicated in transcriptional activation of CDKN1A and the function of the protein involves STAT3 and SRC. And also it is involved in the transcription of rRNA genes by RNA Pol I and enhances protein synthesis and cell growth.