AF/LE Purified Anti-Human/Mouse/Rat CD47 Antibody[MIAP410]


catalog number: E-AB-F10160



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse;Rat
Host	Mouse
Isotype	Mouse IgGl, ĸ
Clone	MIAP410
Conjugation	None (AF/LE)
buffer	Sterile PBS, pH 7.2. \leq 1.0 EU per mg of the antibody as determined by the LAL method

Applications	Recommended Dilution	
FCM	$2 \ \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$	
Data		



Rat splenocytes were stained with 0.2 μg AF/LE Purified Anti-Human/Mouse/Rat CD47 Antibody[MIAP410] (Right) and 0.2 μg Mouse IgG1, κ Isotype Control (Left), followed

by FITC-conjugated Goat Anti-Mouse IgG Secondary

Antibody.

Preparation & Storage	
Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /
	thaw cycles. This preparation contains no preservatives, thus it should be handled
	under aseptic conditions.
Shipping	Ice bag

Background

CD47, also known as Integrin-Associated Protein (IAP), is a membrane protein of about 50 kD with an IgV-like extracelluluar domain, a five membrane-spanning segment and a short terminal cytoplasmic region. It is widely expressed on many cell types and often associated with beta 3 integrins. It has been reported that CD47 functions as a self marker. Red cells lacking CD47 were rapidly cleared from the bloodstream by splenic macrophages. By binding to SIRPα, CD47 controls hemostatic innate immune functions, such as phagocytosis and cell trafficking.

For Research Use Only