







COX15 Polyclonal Antibody

catalog number: E-AB-65067

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse;Rat
Immunogen	A synthetic peptide of human COX15 (NP_510870.1).
Host	Rabbit
Isotype	Rabbit IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
IHC	1:50-1:100
IF	1:50-1:100

Data	
	
Immunohistochemistry of paraffin-embedded Rat spleen using COX15 Polyclonal Antibody at dilution of 1:100 (40x lens).	Immunohistochemistry of paraffin-embedded Rat heart using COX15 Polyclonal Antibody at dilution of 1:100 (40x lens).
	
Immunohistochemistry of paraffin-embedded Human lung cancer using COX15 Polyclonal Antibody at dilution of 1:100 (40x lens).	Immunohistochemistry of paraffin-embedded Human thyroid cancer using COX15 Polyclonal Antibody at dilution of 1:100 (40x lens).
	
Immunohistochemistry of paraffin-embedded Human placenta using COX15 Polyclonal Antibody at dilution of 1:100 (40x lens).	Immunohistochemistry of paraffin-embedded Mouse spleen using COX15 Polyclonal Antibody at dilution of 1:100 (40x lens).


For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.7



Immunofluorescence analysis of L929 cells using COX15 Polyclonal Antibody at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

Cytochrome c oxidase (COX), the terminal component of the mitochondrial respiratory chain, catalyzes the electron transfer from reduced cytochrome c to oxygen. This component is a heteromeric complex consisting of 3 catalytic subunits encoded by mitochondrial genes and multiple structural subunits encoded by nuclear genes. The mitochondrially-encoded subunits function in electron transfer, and the nuclear-encoded subunits may function in the regulation and assembly of the complex. This nuclear gene encodes a protein which is not a structural subunit, but may be essential for the biogenesis of COX formation and may function in the hydroxylation of heme O, according to the yeast mutant studies. This protein is predicted to contain 5 transmembrane domains localized in the mitochondrial inner membrane. Alternative splicing of this gene generates two transcript variants diverging in the 3' region.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.7