

PE/Cyanine5.5 Anti-Human CD70 Antibody[113-16]

Catalog Number: AN00424I

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity	Human
Host	Mouse
Isotype	Mouse IgG1, κ
Clone No.	113-16
Isotype Control	PE/Cyanine5.5 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792I]
Conjugation	PE/Cyanine 5.5
Conjugation Information	PE/Cyanine 5.5 is designed to be excited by the Blue (488 nm), Green (532 nm) and yellow-green (561 nm) lasers and detected using an optical filter centered near 690 nm (e.g., a 690/50 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.


Applications

Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 µL staining volume or per 100 µL of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of the U266 cell line with PE/Cyanine5.5 Anti-Human CD70 Antibody[113-16] (filled gray histogram) or PE/Cyanine5.5 Mouse IgG1, κ Isotype Control (empty black histogram). Total viable cells were used for analysis.

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	GPIX;Glycoprotein IX;GP9;Glycoprotein 9
Uniprot ID	P32970
Gene ID	970

For Research Use Only

Background

CD70, also known as CD27L, is a 50 kD type II transmembrane glycoprotein and member of the tumor necrosis factor superfamily. CD70 is expressed on activated T, B and NK cells, activated plasmacytoid dendritic cells (pDCs), and chronic B cell lymphocytic leukemia and large B cell lymphomas. CD70 costimulates T cell proliferation and differentiation. It plays a role in the pDC-induced B cell differentiation. The ligand of CD70 is CD27.