Recombinant Mouse CD160/BY55 Protein (His Tag)

Catalog Number: PKSM041353

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Species	Mouse	
Source	HEK293 Cells-derived Mouse CD160/BY55 protein Gly28-Ser160, with an C-terminal	
	His	
Calculated MW	15.8 kDa	
Observed MW	20-25 kDa	
Accession	Q8VC80	
Bio-activity	Not validated for activity	
Properties		
Purity	> 95 % as determined by reducing SDS-PAGE.	
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.	
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80	
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of	
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.	
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants	
	before lyophilization.	
	Please refer to the specific buffer information in the printed manual.	
Reconstitution	Please refer to the printed manual for detailed information.	



kDa 170 130 95 72 55 43 34 26	MK	R
17	1	
10	eni	

Background

CD160 antigen is a cell membrane protein which contains one Ig-likeV-type (immunoglobulin-like) domain. CD160 is a GP I-anchored lymphocyte surface receptor in which expression is mostly restricted to the highly cytotoxic CD56(dim) CD1 6(+) peripheral blood NK subset. CD160 is a receptor showing broad specificity for both classical and non-classical MHC class I molecules. CD160 is expressed in spleen, peripheral blood, and smal lintestine. Expression of CD160 is restricted to functional NK and T cytotoxic lymphocytes. CD160 acts as a co-activator receptor for CD3-induced proliferation of CD4+ CD160+ T cells isolated from inflammatory skin lesions. Activated NK lymphocytes release a soluble form of CD160 that functionally impairs the MHC-I-specific cytotoxic CD8(+) T lymphocyte responsiveness.

For Research Use Only

Tel:400-999-2100