# **Elabscience**®

## ALIX / PDCD6IP Polyclonal Antibody

#### catalog number: E-AB-68349


Note: Centrifuge before opening to ensure complete recovery of vial contents.

1:50-1:200

Description	
Reactivity	Human;Mouse;Rat
Immunogen	Recombinant fusion protein of human ALIX / PDCD6IP
Host	Rabbit
Is otype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
WB	1:200-1:2000

IF


#### Data



a ged a

Western blot analysis of extracts of various cell lines using ALIX / PDCD6IP Polyclonal Antibody at 1:1000 dilution.

### Observed-MW:105 kDa Calculated-MW:30 kDa/96 kDa



Immunofluorescence analysis of L929 cells using ALIX / PDCD6IP Polyclonal Antibody at dilution of 1:100. Blue: DAPI for nuclear staining.

Immunofluorescence analysis of U-251MG cells using ALIX

/ PDCD6IP Polyclonal antibody at dilution of 1:100. Blue:

#### DAPI for nuclear staining.

Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

#### Background

### For Research Use Only

# **Elabscience**®

This gene encodes a protein that functions within the ESCRT pathway in the abscission stage of cytokinesis, in intralumenal endosomal vesicle formation, and in enveloped virus budding. Studies using mouse cells have shown that overexpression of this protein can block apoptosis. In addition, the product of this gene binds to the product of the PDCD6 gene, a protein required for apoptosis, in a calcium-dependent manner. This gene product also binds to endophilins, proteins that regulate membrane shape during endocytosis. Overexpression of this gene product and endophilins results in cytoplasmic vacuolization, which may be partly responsible for the protection against cell death. Several alternatively spliced transcript variants encoding different isoforms have been found for this gene. Related pseudogenes have been identified on chromosome 15.

#### For Research Use Only