

A Reliable Research Partner in Life Science and Medicine

# Recombinant Rat 4-1BB/TNFRSF9/CD137 Protein(Fc Tag)

Catalog Number: PDMR100066

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Species Rat

Source HEK293 Cells-derived Rat 4-1BB/TNFRSF9/CD137 proteins Thr24-Val188,with an

C-terminal Fc

 Calculated MW
 43.0 kDa

 Observed MW
 55-60 kDa

 Accession
 Q4V895

Bio-activity Not validated for activity

#### **Properties**

**Purity** > 90% as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU/mg of the protein as determined by the LAL method

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -

80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping

This product is provided as lyophilized powder which is shipped with ice packs.

Formulation


Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5%

Mannitol.

Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution

of 0.5 mg/mL. Concentration is measured by UV-Vis.

### Data



SDS-PAGE analysis of Rat 4-1BB/TNFRSF9/CD137 proteins , 2µg/lane of Recombinant Rat 4-1BB/TNFRSF9/CD137 proteins was resolved with SDS-PAGE under reducing conditions , showing bands at 55-60 KD

## **Background**

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com

Rev. V1.8

# Elabscience®

#### Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

CD137 (also known as 4-1BB) is a surface co-stimulatory glycoprotein originally described as present on activated T lymphocytes , which belongs to the tumor necrosis factor (TNF) receptor superfamily. It is expressed mainly on activated CD4+ and CD8+ T cells , and binds to a high-affinity ligand (4-1BBL) expressed on several antigen-presenting cells such as macrophages and activated B cells. Upon ligand binding , 4-1BB is associated with the tumor necrosis factor receptor—associated factors (TRAFs) , the adaptor protein which mediates downstream signaling events including the activation of NF-kappaB and cytokine production. 4-1BB signaling either by binding to 4-1BBL or by antibody ligation delivers signals for T-cell activation and growth , as well as monocyte proliferation and B-cell survival , and plays an important role in the amplification of T cell-mediated immune responses. In addition , CD137 and CD137L are expressed in different Human primary tumor tissues , suggesting that they may influence the progression of tumors. Crosslinking of CD137 on activated T cells has shown promise in enhancing anti-tumor immune responses in murine models , and agonistic anti-CD137 antibodies are currently being tested in phase I clinical trials. Soluble forms of CD137 (sCD137) are generated by differential splicing. sCD137 can bind to CD137 ligand to antagonize the costimulatory activities of the membrane-bound CD137 and reduce T cell proliferation and IL-2 secretion.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com