Recombinant Human Annexin A6/ANXA6 Protein (His Tag)

Catalog Number: PKSH032076

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | | |
|----------------|-----|--|
| Species | | Human |
| Source | | E.coli-derived Human Annexin A6; ANXA6 protein Ala2-Asp673, with an C-terminal |
| Source | | His |
| Calculated MW | | 76.9 kDa |
| Observed MW | | 60-90 kDa |
| Accession | | P08133 |
| Bio-activity | | Not validated for activity |
| Properties | | |
| Purity | | > 95 % as determined by reducing SDS-PAGE. |
| Endotoxin | | < 1.0 EU per µg of the protein as determined by the LAL method. |
| Storage | | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 |
| | | °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of |
| | | reconstituted samples are stable at $< -20^{\circ}C$ for 3 months. |
| Shipping | | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | | Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. |
| | | Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants |
| | | before lyophilization. |
| | | Please refer to the specific buffer information in the printed manual. |
| Reconstitution | | Please refer to the printed manual for detailed information. |
| Data | | |
| | kDa | MK R |
| | 120 | |
| | 90 | |
| | 60 | |
| | | |
| | 40 | |
| | 30 | |
| | 50 | |

> 95 % as determined by reducing SDS-PAGE.

20

Background

Annexin A6 (ANAX6) belongs to a family of calcium-dependent membrane and phospholipid binding proteins. Annexin A6 is a secreted protein and locates on the cell surface. Annexin A6 contains 8 annexin repeats separated by linking sequences of variable lengths. A pair of annexin repeats may form one binding site for calcium and phospholipid. ANXA6 is involved in the regulation of the release of Ca2+ from intracellular stores and may be associated with CD21. In addition, ANXA6 has been implicated in mediating the endosome aggregation and vesicle fusion in secreting epithelia during exocytosis.