

FGFR1OP2 Polyclonal Antibody

catalog number: E-AB-15033

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human;Mouse;Rat
Immunogen	Recombinant protein of human FGFR1OP2
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.


Applications Recommended Dilution

IHC	1:50-1:200
------------	------------

Data



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using FGFR1OP2 Polyclonal Antibody at dilution 1:40



Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using FGFR1OP2 Polyclonal Antibody at dilution 1:40

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

Acidic and basic fibroblast growth factors (FGFs) are members of a family of multifunctional polypeptide growth factors that stimulate proliferation of cells of mesenchymal, epithelial and neuroectodermal origin. Like other growth factors, FGFs act by binding and activating specific cell surface receptors which include the Flg receptor (FGFR-1) and the Bek receptor (FGFR-2), as well as FGFR-3, FGFR-4, FGFR-5 and FGFR-6. FGFR1OP2 (FGFR1 oncogene partner 2), also known as HSPC123, is a 253 amino acid cytoplasmic protein that is expressed in spleen, thymus and bone marrow and is involved in wound healing under normal cellular conditions. Additionally, FGFR1OP2 may also exist as an aberrant fusion protein with Flg and it is thought that the FGFR1OP2-Flg mutant may play a role in the pathogenesis of stem cell myeloproliferative disorder (MPD). Multiple isoforms of FGFR1OP2 exist due to alternative splicing events.

For Research Use Only