

## Recombinant Human CLC Protein(GST Tag)

**Catalog Number:** PDEH100645

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	E.coli-derived Human CLC protein Ser2-Arg142, with an N-terminal GST
<b>Calculated MW</b>	41.5 kDa
<b>Observed MW</b>	36 kDa
<b>Accession</b>	Q05315
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95% as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 10 EU/mg of the protein as determined by the LAL method
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
<b>Reconstitution</b>	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

### Data



SDS-PAGE analysis of Human CLC proteins, 2µg/lane of Recombinant Human CLC proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 36 KD

### Background

Cardiotrophin-like cytokine (CLC), also referred to as novel neurotrophin-1 (NNT-1) or B cell-stimulating factor-3 (BSF-3), is a new member of the IL-6 family of structurally related cytokines that includes IL-6, CNTF, LIF, CT-1, IL-11 and OSM. All family members share the receptor subunit gp130 that belong to the type I cytokine receptor superfamily. Ligand binding leads to gp130 homodimerization or heterodimerization (with LIF receptor or OSM receptor beta ), and induces cell signaling and functional activity. For several family members, including CNTF, IL-6, and IL-11, binding of the ligand to a specific receptor alpha subunit (CNTF R alpha, IL-6 R alpha, or IL-11 R alpha ) is required prior to gp130 homo- or hetero-dimerization.

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017