



A Reliable Research Partner in Life Science and Medicine

Human BCAM Antibody Pair Set

Catalog No. E-KAB-0697 Applications ELISA

Synonyms CD239;BCAM;AU;LU;MSK19;Lutheran Blood Group Glycoprotein;Auberger B antigen;B-

CAM cell surface glycoprotein

Kit components & Storage

Title	Specifications	Storage
Human BCAM Capture Antibody	1 vial, 100 μ g	Store at -20℃. Avoid freeze / thaw
		cycles.
Human BCAM Detection Antibody	1 vial, 50 μL	Store at -20°C. Avoid freeze / thaw
(Biotin)		cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0697)		
		Human BCAM Capture Antibody	Human BCAM Detection Antibody	
			(Biotin)	
Immunogen	Immunogen	Recombinant Human BCAM protein	Recombinant Human BCAM protein	
Information	Swissprot	P50895		
Product details	Reactivity	Human	Human	
	Host	Rabbit	Rabbit	
	Conjugation	Unconjugated	Biotin	
	Concentration	0.5mg/mL	/	
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%	
		glycerol, pH 7.4	protective protein, 50% glycerol, pH	
			7.4	
	Purify	Affinity purification	Affinity purification	
	Specificity	Detects Human BCAM in ELISAs.		

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com



A Reliable Research Partner in Life Science and Medicine

Applications

Human BCAM Sandwich ELISA Assay

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4ug/mL	Human BCAM Capture	
Capture		Antibody	10
			8 11
			Optical Density
ELISA	1:1000-1:10000	Human BCAM Detection	0 O O O O O
Detection		Antibody (Biotin)	•
			0.01
			Human BCAM Concentration(pg/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

Human CD239 is a type I transmembrane glycoprotein that is encoded by B-CAM which belongs to the Ig gene superfamily. Full-length BCAM (CD239) is comprised of an extracellular region with two N-terminal V-type Ig-like domains followed by three C2-type Ig-like domains, a transmembrane region, and cytoplasmic domain of 59 amino acid residues. CD239 is expressed on erythrocytes in the peripheral blood and on the basal layer of vascular endothelial cells or some epithelial cells.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com