



SNAPC5 Polyclonal Antibody

catalog number: E-AB-18025

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|--------------|--|
| Reactivity | Human;Mouse |
| Immunogen | Synthetic peptide of human SNAPC5 |
| Host | Rabbit |
| Isotype | IgG |
| Purification | Antigen affinity purification |
| Conjugation | Unconjugated |
| Buffer | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |
| Applications | Recommended Dilution |
| IHC | 1:30-1:150 |

| Data | |
|--|---|
|  |  |
| Immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using SNAPC5 Polyclonal Antibody at dilution of 1:25(×200) | Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using SNAPC5 Polyclonal Antibody at dilution of 1:25(×200) |

| Preparation & Storage | |
|-----------------------|--|
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |

| Background | |
|---|--|
| <p>This gene encodes a subunit of the small nuclear RNA (snRNA)-activating protein complex that plays a role in the transcription of snRNA genes. This complex binds to the promoters of snRNA genes transcribed by either RNA polymerase II or III and recruits other regulatory factors to activate snRNA gene transcription. The encoded protein may play a role in stabilizing this complex. A pseudogene of this gene has been identified on chromosome 6.</p> | |