Elabscience®

Elab Fluor[®] 700 Anti-Mouse CD18 Antibody[M18/2]


Catalog Number: E-AB-F1113UM1

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Mouse
Host	Rat
Isotype	Rat lgG2a, κ
Clone No.	M18/2
Isotype Control	Elab Fluor [®] 700 Rat IgG2a, к Isotype Control[2А3] [Product E-AB-F09833М1]
Conjugation	Elab Fluor [®] 700
Conjugation Information	Elab Fluor [®] 700 is designed to be excited by the Red laser (627-640 nm) and detected using an optical filter centered near 719 nm (e.g., a 725/40 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the

reagent to obtain optimal results [The recommended concentration is 0.1-1 μ g/10⁶ cells in 100 μ L volume].

Data



Staining of C57BL/6 murine splenocytes with Elab Fluor[®] 700 Anti-Mouse CD18 Antibody[M18/2] (filled gray

histogram) or Elab Fluor[®] 700 Rat IgG2a, κ Isotype Control (empty black histogram). Total viable cells were used for analysis.

d from prolonged
d from prolonged
bunit beta;
1

For Research Use Only

Elabscience®

Gene ID Background

16414

CD18 is a 95 kD protein, also known as integrin β 2 subunit. It is expressed on all leukocytes. CD18, in association with integrin α chain CD11a, CD11b, and CD11c forms LFA-1, Mac-1, and α X β 2, respectively, and plays an important role in leukocytes adhesion. The CD18 integrin complexes bind ICAM-1 (CD54), ICAM-2 (CD102), ICAM-3 (CD50), iC3b, and fibrinogen.