SARS-COV-2 Spike RBD Polyclonal Antibody (2019-nCoV)

catalog number: E-AB-V1006

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Note: Centrifuge before opening to ensure complete recovery of vial contents:	
Description	
Reactivity	SARS-COV2
Immunogen	Recombinant 2019-nCoV Spike Protein (RBD)
Host	Rabbit
Isotype	IgG
Buffer	0.2 µm filtered solution in PBS.
Applications	Recommended Dilution
WB	1:1000-1:5000
Data	
	12 13 14 16
Western Blot analysis of SARS-CoV2-Spike protein using	
SARS-COV/SARS-COV-2 Spike RBD Polyclonal	
Antibody(2019-nCoV)at dilution of 1:2000	
Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the
	temperature recommended.
Background	

Background

Protein S (PROS1) is glycoprotein and expressed in many cell types supporting its reported involvement in multiple biological processes that include coagulation, apoptosis, cancer development and progression, and the innate immune response. Known receptors bind S1 are ACE2, angiotensin-converting enzyme 2, DPP4, CEACAM etc. The spike (S) glycoprotein of coronaviruses is known to be essential in the binding of the virus to the host cell at the advent of the infection process. Most notable is severe acute respiratory syndrome (SARS). The severe acute respiratory syndrome-coronavirus (SARS-CoV) spike (S) glycoprotein alone can mediate the membrane fusion required for virus entry and cell fusion. It is also a major immunogen and a target for entry inhibitors. It's been reported that 2019-nCoV can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.