Elabscience Biotechnology Co., Ltd.



A Reliable Research Partner in Life Science and Medicine

CCNE1 Polyclonal Antibody

catalog number: E-AB-19281

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Synthetic peptide of human CCNE1

Host Rabbit Isotype IgG

Purification Antigen affinity purification

Conjugation Unconjugated


Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications Recommended Dilution

IHC 1:30-1:150

Data







Immunohistochemistry of paraffin-embedded Human cervical cancer tissue using CCNE1 Polyclonal Antibody at dilution of 1:25(×200)

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK2, whose activity is required for cell cycle GI/S transition. This protein accumulates at the GI-S phase boundary and is degraded as cells progress through S phase. Overexpression of this gene has been observed in many tumors, which results in chromosome instability, and thus may contribute to tumorigenesis. This protein was found to associate with, and be involved in, the phosphorylation of NPAT protein (nuclear protein mapped to the ATM locus), which participates in cel l-cycle regulated histone gene expression and plays a critical role in promoting cell-cycle progression in the absence of pRB.

For Research Use Only