


IDO1 Polyclonal Antibody

catalog number: E-AB-93146

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description	
Reactivity	Human;Rat
Immunogen	Recombinant fusion protein of human IDO1
Host	Rabbit
Isotype	Rabbit IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
WB	1:500-1:2000

Data



Western blot analysis of extracts of SKOV3 cells using IDO1 Polyclonal Antibody at 1:500 dilution.

Observed-MW:40 kDa
Calculated-MW:45 kDa



Western blot analysis of extracts of A-549 cells using IDO1 Polyclonal Antibody at 1:500 dilution. A-549 cells were treated by hIFN- γ (100 ng/mL) at 37°C for 48 hours.

Observed-MW:40 kDa
Calculated-MW:45 kDa

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

This gene encodes indoleamine 2,3-dioxygenase (IDO) - a heme enzyme that catalyzes the first and rate-limiting step in tryptophan catabolism to N-formyl-kynurenine. This enzyme acts on multiple tryptophan substrates including D-tryptophan, L-tryptophan, 5-hydroxy-tryptophan, tryptamine, and serotonin. This enzyme is thought to play a role in a variety of pathophysiological processes such as antimicrobial and antitumor defense, neuropathology, immunoregulation, and antioxidant activity. Through its expression in dendritic cells, monocytes, and macrophages this enzyme modulates T-cell behavior by its peri-cellular catabolization of the essential amino acid tryptophan.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.8