Recombinant Human NEDD8

Catalog Number: PKSH034041

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | | | |
|----------------|------------------------------------------------------------------------------------------|--|--|
| Species | Human | | |
| Source | E.coli-derived Human NEDD8 protein Met1-Gly76 | | |
| Calculated MW | 8.6 kDa | | |
| Observed MW | 9 kDa | | |
| Accession | Q15843 | | |
| Bio-activity | Not validated for activity | | |
| Properties | | | |
| Purity | >95 % as determined by reducing SDS-PAGE. | | |
| Endotoxin | < 1.0 EU per µg of the protein as determined by the LAL method. | | |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 | | |
| | °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of | | |
| | reconstituted samples are stable at $< -20^{\circ}$ C for 3 months. | | |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. | | |
| Formulation | Den Lyophilized from a 0.2 μm filtered solution of 20mM Tris-HCl, 250mM NaCl, pH | | |
| | Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants | | |
| | before lyophilization. | | |
| | Please refer to the specific buffer information in the printed manual. | | |
| Reconstitution | Please refer to the printed manual for detailed information. | | |

Data

| kDa | МК | R | |
|-----------------|----|---|--|
| 120 90 60 | | | |
| 40 | | | |
| 30 | - | | |
| 20 | | | |
| 14 | - | - | |
| | | | |

> 95 % as determined by reducing SDS-PAGE.

Background

Human NEDD8 shares 60% amino acid sequence identity to ubiquitin. The only known substrates of NEDD8 modification are the cullin subunits of SCF ubiquitin E3 ligases. The NEDDylation of cullins is critical for the recruitment of E2 to the ligase complex, thus facilitating ubiquitin conjugation. NEDD8 modification has therefore been implicated in cell cycle progression and cytoskeletal regulation.