

## Recombinant Human CD47/IAP Protein (Fc & Avi Tag)

**Catalog Number:** PKSH033799

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human CD47/IAP protein Gln19-Pro139, with an C-terminal Fc & Avi
<b>Calculated MW</b>	42.7 kDa
<b>Observed MW</b>	60-70 kDa
<b>Accession</b>	Q08722
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
	Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

CD47(Integrin-Associated Protein;IAP) is a 40 - 60 kDa variably glycosylated atypical member of the immunoglobulin superfamily. The ubiquitously expressed CD47 binds to SIRP family members on macrophages; neutrophils; and T cells. CD47 is involved in the increase in intracellular calcium concentration that occurs upon cell adhesion to extracellular matrix. The protein is also a receptor for the C-terminal cell-binding domain of thrombospondin; and it may play a role in membrane transport and signal transduction. This protein has broad tissue distribution; and is reduced in expression on Rh erythrocytes.

### For Research Use Only