A Reliable Research Partner in Life Science and Medicine

# VE-Cadherin Polyclonal Antibody(Capture/Detector)

catalog number: AN000140P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description

Reactivity Human

Immunogen Recombinant Human VE-Cadherin protein expressed by E.coli

Host Rabbit
Isotype Rabbit IgG


**Purification** Antigen Affinity Purification

**Buffer** Phosphate buffered solution, pH 7.2, containing 0.05% proclin 300.

## **Applications** Recommended Dilution

ELISA Capture 2-8 μg/mL ELISA Detector 0.1-0.4 μg/mL

#### Data



Sandwich ELISA-Recombinant Human VE-Cadherin protein standard curve. Background subtracted standard curve using VE-Cadherin antibody(AN000140P)(Capture), VE-Cadherin antibody(AN000140P)(Detector) in sandwich ELISA. The reference range value for Recombinant Human VE-Cadherin protein is 31.25-2000 pg/mL.

## **Preparation & Storage**

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

**Shipping** The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

## Background

Cadherins are calcium-dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells, cadherins may thus contribute to the sorting of heterogeneous cell types. This cadherin may play a important role in endothelial cell biology through control of the cohesion and organization of the intercellular junctions. It associates with alpha-catenin forming a link to the cytoskeleton. Acts in concert with KRIT1 and PAIS1 to establish and maintain correct endothelial cell polarity and vascular lumen. These effects are mediated by recruitment and activation of the Par polarity complex and RAP1B. Required for activation of PRKCZ and for the localization of phosphorylated PRKCZ, PARD3, TIAM1 and RAP1B to the cell junction.

#### For Research Use Only

Fax: 1-832-243-6017