

A Reliable Research Partner in Life Science and Medicine

Recombinant Calreticulin Monoclonal Antibody

catalog number: AN301032L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse; Rat

Immunogen Recombinant Human Calreticulin protein


HostRabbitIsotypeIgG,κCloneB783PurificationProtein A

Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications Recommended Dilution

IHC 1:200-1:1000 **WB** 1:1000-1:5000

Data




Western Blot with Recombinant Calreticulin Monoclonal Antibody at dilution of 1:1000 dilution. Lane A: NIH-3T3 cells.

Immunohistochemistry of paraffin-embedded human kidney tissue using Recombinant Calreticulin Monoclonal Antibody at dilution of 1:200.

Rev. V1.1

Observed-MW:55 kDa Calculated-MW:48 kDa



Immunohistochemistry of paraffin-embedded rat stomach using Recombinant Calreticulin Monoclonal Antibody at dilution of 1:200.

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping lce bag

Background

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Calreticulin is a multifunctional protein that acts as a major Ca(2+)-binding (storage) protein in the lumen of the endoplasmic reticulum. It is also found in the nucleus, suggesting that it may have a role in transcription regulation. Calreticulin binds to the synthetic peptide KLGFFKR, which is almost identical to an amino acid sequence in the DNA-binding domain of the superfamily of nuclear receptors. Calreticulin binds to antibodies in certain sera of systemic lupus and Sjogren patients which contain anti-Ro/SSA antibodies, it is highly conserved among species, and it is located in the endoplasmic and sarcoplasmic reticulum where it may bind calcium. The amino terminus of calreticulin interacts with the DNA-binding domain of the glucocorticoid receptor and prevents the receptor from binding to its specific glucocorticoid response element.

For Research Use Only