



A Reliable Research Partner in Life Science and Medicine

APC Anti-Human CD50 Antibody[CBR-IC3/1]

Catalog Number: AN00318E

Note: Centrifuge before opening to ensure complete recovery of vial contents.

_		400	
Desc		7 77	٦n
レヒコレ	ш		711

Reactivity Human Host Mouse

Isotype Mouse IgG1, κ **Clone No.** CBR-IC3/1

Isotype Control APC Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792E]

Conjugation APC

Conjugation Information APC is designed to be excited by the Red (627-640 nm) laser and detected using an

optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.

Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names ICAM-3
Uniprot ID P32942
Gene ID 3385

Background CD50 is a 120-140 kD type I membrane protein also known as intercellular adhesion

Web: www.elabscience.cn

molecule 3 (ICAM-3). It is broadly expressed on leukocytes, endothelial cells, and Langerhans cells. CD50 is a member of the lg superfamily. It provides costimulatory signal for T cells and regulates cell adhesion through interaction with LFA-1 (CD11a/

CD18, $\alpha L/\beta 2$) integrin.