

Recombinant Human LIF protein (His tag)

Catalog Number:PDMH100067



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms	Leukemia Inhibitory Factor;LIF;Differentiation-Stimulating Factor;D Factor;Melanoma-Derived LPL Inhibitor;MLPLI;Emfifermin;LIF;HILDA;CDF;DIA
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Phe202
Accession	P15018
Calculated Molecular Weight	19.8 kDa
Observed molecular weight	38 kDa
Tag	C-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Background

Leukemia inhibitory factor (LIF) is a pleiotropic glycoprotein belonging to the IL-6 family of cytokines. It's involved in growth promotion and cell differentiation of different types of target cells; influence on bone metabolism; cachexia; neural development; embryogenesis and inflammation. LIF has potent proinflammatory property; being the inducer of the acute phase protein synthesis and affecting the cell recruitment into the area of damage or inflammation. LIF is also one of the cytokines that are capable to regulate the differentiation of embryonic stem cells; hematopoietic and neuronal cells. LIF binds to the specific LIF receptor (LIFR- α) which forms a heterodimer with a specific subunit common to all members of that family of receptors; the GP130 signal transducing subunit. This leads to activation of the JAK/STAT and MAPK cascades. Due to its polyfunctional activities; LIF is involved in the pathogenic events and development of many diseases of various origin.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017