

## Recombinant Rat IL2/IL-2/Interleukin-2 Protein (His Tag)

**Catalog Number:** PDER100199

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Rat
<b>Source</b>	E.coli-derived Rat IL2 protein Ala21-Gln155, with an C-terminal His
<b>Calculated MW</b>	14.7 kDa
<b>Observed MW</b>	15 kDa
<b>Accession</b>	P17108
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95% as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 10 EU/mg of the protein as determined by the LAL method
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
<b>Reconstitution</b>	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

### Data



SDS-PAGE analysis of Rat IL2/IL-2/Interleukin-2 proteins, 2 µg/lane of Recombinant Rat IL2/IL-2/Interleukin-2 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 15 kDa.

### Background

Interleukin-2(IL-2)is a O-glycosylated four α-helix bundle cytokine that has potent stimulatory activity for antigenactivated T cells. It is expressed by CD4+ and CD8+ T cells, γδ T cells, B cells, dendritic cells, and eosinophils. Mature rat IL-2 shares 66% and 73% amino acid sequence identity with human and mouse IL-2,respectively. The receptor for IL-2 consists of three subunits that are present on the cell surface in varying preformed complexes. IL-2 is a powerful immunoregulatory lymphokine produced by T-cells in response to antigenic or mitogenic stimulation. IL-2/IL-2R signaling is required for T-cell proliferation and other fundamental functions that are essential for the immune response. IL-2 stimulates growth and differentiation of B-cells, NK cells, lymphokine-activated killer cells, monocytes, macrophages and oligodendrocytes.

### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017