Elabscience®

Flag-Tag Monoclonal Antibody


catalog number: E-AB-48025

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	All
Immunogen	Synthetic peptide corresponding to Flag tag conjugated to keyhole limpet
	haemocyanin.
Host	Mouse
Is otype	IgG2a
Clone	3H9
Purification	Protein A/G Purification
Buffer	PBS, pH 7.2. Contains 0.05% Proclin300.
Applications	Recommended Dilution
WB	1:5000-10000
IF	1:500-2000
IP	3µg/sample
FCM	0.2µg
Data	
	55KD
	43KD FLAG
	34KD

Western blotting with anti-FLAG monoclonal antibody at

26KD 17KD




dilution of 1:5000.lane 1: HEK 293 whole cell lysate, Lane 2: the FLAG-GFP, using anti-Flag-Tag Monoclonal Antibody

FLAG tag transfected HEK 293 cell lysates Observed-MW:45 kDa Calculated-MW:45 kDa



at 1:2000 dilution.



Lane 1:HEK-293 transfected with DDDDK-tagged (input).Lane 2: Mouse monoclonal IgG2a instead of E-AB-48025 in HEK-293T transfected with DDDDK-tagged.Lane 3: E-AB-48025 IP in HEK-293T transfected with DDDDK- Control stained with 0.2ug mouse IgG and AF647 conjugated

tagged.

1x10⁶ CHO cells Transfected with a Flag plasmid were stained with 0.2ug Anti-Flag tag mAb(E-AB-48025) and AF647 conjugated Goat Anti-mouse IgG (H+L);lsotype

Goat Anti-mouse IgG (H+L)

Preparation & Storage		
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.	
For Research Use Only		
Toll-free: 1-888-852-8623	Tel: 1-832-243-6086	Fax: 1-832-243-6017
Web:www.elabscience.com	Email:techsupport@elabscience.com	

Elabscience®

Shipping

The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended.

Background

FLAG-tag is a polypeptide protein tag that can be added to a protein using recombinant DNA technology, having the sequence motif DYKDDDDK. It has been used for studying proteins in living cells and for protein purification by affinity chromatography. This antibody detects FLAG tag in most applications.