Elabscience®

MGAT4B Polyclonal Antibody

catalog number: E-AB-19215

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | | |
|----------------------------------|------------------------------------|--|
| Reactivity | Human;Mouse | |
| Immunogen | Fusion protein of human M | GAT4B |
| Host | Rabbit | |
| Isotype | IgG | |
| Purification | Antigen affinity purification | 1 |
| Conjugation | Unconjugated | |
| Buffer | Phosphate buffered solution | n, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |
| Applications | Recommended Dilution | |
| IHC | 1:50-1:100 | |
| Data | | |
| Immunohistochemistry o | f paraffin-embedded Human cervical | Immunohistochemistry of paraffin-embedded Human |
| | | colorectal cancer tissue using MGAT4B Polyclonal |
| of 1:50(×200) | | Antibody at dilution of 1:50(×200) |
| Preparation & Storage | | |
| Preparation & Storage Storage | | oonths. A void freeze / thaw cycles |

| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. | |
|----------|---|--|
| Shipping | The product is shipped with ice pack, upon receipt, store it immediately at the | |
| | temperature recommended. | |

Background

This gene encodes a key glycosyltransferase that regulates the formation of tri- and multiantennary branching structures in the Golgi apparatus. The encoded protein, in addition to the related isoenzyme A, catalyzes the transfer of N-acetylglucosamine (GlcNAc) from UDP-GlcNAc in a beta-1,4 linkage to the Man-alpha-1,3-Man-beta-1,4-GlcNAc arm of R-Man-alpha-1,6(GlcNAc-beta-1,2-Man-alpha-1,3)Man-beta-1,4-GlcNAc-beta-1,4-GlcNAc-beta-1-Asn. The encoded protein may play a role in regulating the availability of serum glycoproteins, oncogenesis, and differentiation.