


GAPDH Monoclonal Antibody

catalog number: E-AB-48016

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description	
Reactivity	Human;Mouse;Rat ;Bovine;Caprine;Guinea pig;Rabbit;Porcine;
Immunogen	Recombinant human GAPDH protein expressed by E.coli
Host	Mouse
Isotype	IgG1
Clone	3B3
Purification	Protein A/G Purification
Conjugation	Unconjugated
Buffer	PBS with 0.05% proclin 300, 1% protective protein and 50% glycerol,pH7.4
Applications	
WB	1:5000-10000

Data




Western Blot analysis of GAPDH in various cell lines at dilution of 1:5000. Lane1: DU145 whole cell lysate, Lane2: OS-RC-2 whole cell lysate, Lane3:T-47D whole cell Lysates, Lane4: HEC-1B whole cell lysate, Lane5: HepG2 whole cell lysate, Lane6: HEK-293 whole cell lysate, Lysates/proteins at 20g per lane.

Observed-MW:36 kDa
Calculated-MW:36 kDa



Western blot analysis of GAPDH in various tissues of different species at dilution of 1:5000. Lane1 : Rat heart, whole tissue lysate, Lane2: Porcine brain, whole tissue lysate, Lane3: Bovine kidney, whole tissue lysate, Lane4: Guinea pig liver, whole tissue lysate, Lane5: Rabbit pancreas, whole tissue ysate, Lane6: Caprine spleen, whole tissue lysate. Lysates/proteins at 20g per lane.

Observed-MW:36 kDa
Calculated-MW:36 kDa



Immunofluorescent analysis of 293F cells using anti-GAPDH monoclonal antibody at 1:400 dilution.

Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended.

For Research Use Only

Background

GAPDH is an enzyme of 37kDa that catalyzes the sixth step of glycolysis and thus serves to break down glucose for energy and carbon molecules. Because the GAPDH gene is often stably and constitutively expressed at high levels in most tissues and cells, it is commonly used as a loading control for western blot and as a control for qPCR.