Recombinant Mouse CD40/TNFRSF5 Protein (Fc Tag)

Catalog Number: PKSM041225

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description			
Species	Mouse		
Source	HEK293 Cells-derived Mouse CD40/TNFRSF5 protein Leu20-Arg193, with an C-		
	terminal Fc		
Calculated MW	46.5 kDa		
Observed MW	50-58 kDa		
Accession	P27512		
Bio-activity	Not validated for activity		
Properties			
Purity	>95% as determined by reducing SDS-PAGE.		
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.		
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -8		
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of		
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.		
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.		
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.		
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants		
	before lyophilization.		
	Please refer to the specific buffer information in the printed manual.		
Reconstitution	Please refer to the printed manual for detailed information.		

Data

kDa	MK	R
120		
90 60		22-22-2
40		and the second
30		
20	-	
		1211
14	15-33	Chilling .

Background

Elabscience®

CD40 is a Type I Transmembrane Glycoprotein that belongs to the TNF Receptor Superfamily. CD40 is expressed in B cells, follicular dendritic cells, dendritic cells, activated monocytes, macrophages, endothelial cells, vascular smooth muscle cells, and several tumor cell lines. The extracellular domain of CD40 is characterized by Cysteine rich repeat regions. Interaction of CD40 with its ligand (CD40L) leads to aggregation of CD40 molecules, which in turn interact with cytoplasmic components to initiate signaling pathways. Several different TRAF proteins (adaptor proteins) have been identified to serves as mediators of the signal transduction. CD40 plays an essential role in mediating a broad variety of immune and inflammatory responses including T cell-dependent immunoglobulin class switching, memory B cell development, and germinal center formation.