## **Elabscience**®

### Recombinant Mouse AXL/UFO Protein( Fc Tag)

### Catalog Number: PDMM100113

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description			
Species	Mouse		
Source	Mammalian-derived Mouse AXL/UFO protein Ala19A-Trp445, with an C-terminal Fc		
Calculated MW	71.8 kDa		
Observed MW	90-110 kDa		
Accession	Q00993		
Bio-activity	Not validated for activity		
Properties			
Purity	> 90% as determined by reducing SDS-PAGE.		
Endotoxin	< 1.0 EU/mg of the protein as determined by the LAL method		
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80		
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of		
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.		
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.		
Formulation	Lyophilized from a 0.2 $\mu$ m filtered solution in PBS with 5% Trehalose and 5%		
	Mannitol.		
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of		
	0.5 mg/mL. Concentration is measured by UV-Vis.		

Data

kDa	М	R
80 60	_	1000
40	-	
30	-	
20	-	

SDS-PAGE analysis of Mouse AXL/UFO proteins, 2 µg/lane of Recombinant Mouse AXL/UFO proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 71.8 KD

#### Background

# **Elabscience**®

Axl receptor tyrosine kinase, together with Tyro3 and Mer, constitute the TAM family of receptor tyrosine kinases. In the nervous system, Axl and its ligand Growth-arrest-specific protein 6 (Gas6) are expressed on multiple cell types. Axl functions in dampening the immune response, regulating cytokine secretion, clearing apoptotic cells and debris, and maintaining cell survival. Axl is upregulated in various disease states, such as in the cuprizone toxicity-induced model of demyelination and in multiple sclerosis (MS) lesions, suggesting that it plays a role in disease pathogenesis. Axl expression correlates with poor prognosis in several cancers. Axl mediates multiple oncogenic phenotypes and activation of these RTKs constitutes a mechanism of chemoresistance in a variety of solid tumors. Axl contributes to cell survival, migration, invasion, metastasis and chemosensitivity justify further investigation of Axl as novel therapeutic targets in cancer. The receptor tyrosine kinase AXL is thought to play a role in metastasis. The soluble AXL receptor as a therapeutic candidate agent for treatment of metastatic ovarian cancer. GAS6/AXL targeting as an effective strategy for inhibition of metastatic tumor progression in vivo.