

Anti-2019-nCoV S-mIgG1 Neutralizing Antibody

catalog number: E-AB-V1022

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	SARS-COV2
Immunogen	Recombinant 2019-nCoV S-trimer Protein (His Tag)
Host	Mouse
Isotype	IgG1
Clone	8A5
Conjugation	Unconjugated
buffer	0.2 μm filtered solution in PBS.

Applications


Recommended Dilution

ELISA	1:1000-1:2000
--------------	---------------


Data




Immobilized Recombinant 2019-nCoV S-trimer Protein (C-6His)(Cat#PKSR030489) at 5.0 μg/mL (100 uL/well) can bind Anti-2019-nCoV S-mIgG1 Neutralizing Antibody (8A5), the EC₅₀ is 4.43 ng/mL.



Immobilized Recombinant Recombinant 2019-nCoV S Protein RBD-SD1 (C-6His) (Cat#PKSR030477) at 5.0 ug/mL (100 uL/well) can bind Anti-2019-nCoV S-mIgG1 Neutralizing Antibody (8A5), the EC₅₀ is 6.5 ng/mL.



Anti-2019-nCoV S-mIgG1 Neutralizing Antibody(8A5) can block Human ACE-2 Protein (Avi-His Tag) (Cat#PKSR030493) and 2019-nCoV S-trimer Protein(Cat#PKSR030489) interaction, the IC₅₀ for this effect is 80 ng/mL.



Capture Antibody: E-AB-V1022[8A5]
Detection Antibody: E-AB-V1008[5E8]
ED50: 11.6 μg/mL

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

For Research Use Only

Anti-2019-nCoV S-mIgG1 Neutralizing Antibody

catalog number: E-AB-V1022



Protein S (PROS1) is glycoprotein and expressed in many cell types supporting its reported involvement in multiple biological processes that include coagulation, apoptosis, cancer development and progression, and the innate immune response. Known receptors bind S1 are ACE2, angiotensin-converting enzyme 2, DPP4, CEACAM etc. The spike (S) glycoprotein of coronaviruses is known to be essential in the binding of the virus to the host cell at the advent of the infection process. Most notable is severe acute respiratory syndrome (SARS). The severe acute respiratory syndrome-coronavirus (SARS-CoV) spike (S) glycoprotein alone can mediate the membrane fusion required for virus entry and cell fusion. It is also a major immunogen and a target for entry inhibitors. It's been reported that 2019-nCoV can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine
Tel: 400-999-2100

Email: techsupport@elabscience.cn

Web: www.elabscience.cn

Rev. V1.4