
Recombinant Human Neurotrimin/NTM Protein (His Tag)

Catalog Number: PKSH032802

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| SpeciesHumanSourceHEK293 Cells-derived Human Neurotrimin;NTM protein Gly34-Leu316, terminal HisCalculated MW32.3 kDaObserved MW55-70 kDaAccessionQ9P121-3Bio-activityNot validated for activityPropertiesPurity> 95 % as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU per µg of the protein as determined by the LAL method. | |
|--|-------------------|
| terminal HisCalculated MW32.3 kDaObserved MW55-70 kDaAccessionQ9P121-3Bio-activityNot validated for activityPropertiesPurity> 95 % as determined by reducing SDS-PAGE.Findotoxin< 1.0 EU per µg of the protein as determined by the LAL method.StorageGenerally, lyophilized proteins are stable for up to 12 months when stor °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. A reconstituted samples are stable at < -20°C for 3 months. | |
| Calculated MW32.3 kDaObserved MW55-70 kDaAccessionQ9P121-3Bio-activityNot validated for activityPropertiesPurity> 95 % as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU per µg of the protein as determined by the LAL method. | , with an C- |
| Observed MW55-70 kDaAccessionQ9P121-3Bio-activityNot validated for activityPropertiesPurity> 95 % as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU per μg of the protein as determined by the LAL method. | |
| Accession Q9P121-3 Bio-activity Not validated for activity Properties Purity Purity > 95 % as determined by reducing SDS-PAGE. Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method. Storage Generally, lyophilized proteins are stable for up to 12 months when stor °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. A reconstituted samples are stable at < -20°C for 3 months. | |
| Bio-activityNot validated for activityPropertiesPurity> 95 % as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU per μg of the protein as determined by the LAL method. | |
| Properties Purity > 95 % as determined by reducing SDS-PAGE. Endotoxin < 1.0 EU per μg of the protein as determined by the LAL method. Storage Generally, lyophilized proteins are stable for up to 12 months when stor °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. A reconstituted samples are stable at < -20°C for 3 months. | |
| Purity> 95 % as determined by reducing SDS-PAGE.Endotoxin< 1.0 EU per μg of the protein as determined by the LAL method. | |
| Endotoxin< 1.0 EU per μg of the protein as determined by the LAL method.StorageGenerally, lyophilized proteins are stable for up to 12 months when stor °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. A reconstituted samples are stable at < -20°C for 3 months. | |
| StorageGenerally, lyophilized proteins are stable for up to 12 months when stor °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. A reconstituted samples are stable at < -20°C for 3 months. | |
| °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. A reconstituted samples are stable at < -20 °C for 3 months. | |
| reconstituted samples are stable at $< -20^{\circ}$ C for 3 months. | red at -20 to -80 |
| • | Aliquots of |
| Shipping This product is provided as lyophilized powder which is shipped with it | |
| | ice packs. |
| FormulationLyophilized from a 0.2 μm filtered solution of 20mM PB,150mM NaCl, | pH7.4. |
| Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added a | as protectants |
| before lyophilization. | |
| Please refer to the specific buffer information in the printed manual. | |
| Reconstitution Please refer to the printed manual for detailed information. | |

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Neurotrimin is a glycoprotein. It belongs to the immunoglobulin superfamily and IgLON family. Neurotrimin localizes to the cell membrane and contains three Ig-like C2-type domains. Neurotrimin acts as a glycosylphosphatidylinositol (GPI)anchored neural cell adhesion molecule. Neurotrimin may promote neurite outgrowth and adhesion via homophilic and heterophilic interactions.